



## Photodegradation of amoxicillin by catalyzed $\text{Fe}^{3+}/\text{H}_2\text{O}_2$ process

Xiaoming Li<sup>1,2,3,\*</sup>, Tingting Shen<sup>1,2</sup>, Dongbo Wang<sup>1,2</sup>, Xiu Yue<sup>1,2</sup>, Xian Liu<sup>1,2</sup>, Qi Yang<sup>1,2</sup>,  
Jianbin Cao<sup>1,2</sup>, Wei Zheng<sup>1,2</sup>, Guangming Zeng<sup>1,2</sup>

1. College of Environmental Science and Engineering, Hunan University, Changsha, 410082, China. E-mail: [xmli@hnu.cn](mailto:xmli@hnu.cn)

2. Key Laboratory of Environmental Biology and Pollution Control, Hunan University, Ministry of Education, Changsha 410082, China

3. School of Environment, Guangxi University, Nanning 530004, China

Received 22 February 2011; revised 08 June 2011; accepted 17 June 2011

### Abstract

Three oxidation processes of  $\text{UV-Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  (UV: ultraviolet light; EDTA: ethylenediaminetetraacetic acid),  $\text{UV-Fe}^{3+}/\text{H}_2\text{O}_2$  and  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  were simultaneously investigated for the degradation of amoxicillin at pH 7.0. The results indicated that, 100% amoxicillin degradation and 81.9% chemical oxygen demand ( $\text{COD}_{\text{Cr}}$ ) removal could be achieved in the  $\text{UV-Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  process. The treatment efficiency of amoxicillin and  $\text{COD}_{\text{Cr}}$  removal were found to decrease to 59.0% and 43.0% in the  $\text{UV-Fe}^{3+}/\text{H}_2\text{O}_2$  process; 39.6% and 31.3% in the  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process. Moreover, the results of biodegradability (biological oxygen demand ( $\text{BOD}_5$ )/ $\text{COD}_{\text{Cr}}$  ratio) revealed that the  $\text{UV-Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  process was a promising strategy to degrade amoxicillin as the biodegradability of the effluent was improved to 0.45, compared with the cases of  $\text{UV-Fe}^{3+}/\text{H}_2\text{O}_2$  (0.25) and  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  (0.10) processes. Therefore, it could be deduced that EDTA and UV light performed synergistic catalytic effect on the  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process, enhancing the treatment efficiency. The degradation mechanisms were also investigated via UV-Vis spectra, and high performance liquid chromatography-mass spectra. The degradation pathway of amoxicillin was further proposed.


**Key words:** amoxicillin; catalytic oxidation; photodegradation

**DOI:** 10.1016/S1001-0742(11)60765-1

### Introduction

The occurrence of antibiotics in the aquatic environment has received considerable attention, which are regarded as the potential contaminations due to their continuous input into the ecosystem (Gulkowska et al., 2008; Huber et al., 2003; Kümmerer et al., 2000; Migliore et al., 1997). Another concern about antibiotic residues in the environment is the potential risk for aquatic and terrestrial organisms that exhibit resistance to some antibiotics. It has been reported that bacteria isolated from sewages showed resistance to antibiotics, such as ciprofloxacin, erythromycin and tetracycline (da Silva et al., 2006).

In the past decades, Fenton oxidation or Fenton like processes have been investigated for antibiotics degradation and biodegradability improvement, such as  $\text{Fe}^{2+}/\text{H}_2\text{O}_2$ ,  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$ ,  $\text{Fe}^{2+}/\text{H}_2\text{O}_2/\text{UV}$  and  $\text{Fe}^{3+}/\text{H}_2\text{O}_2/\text{UV}$ . Compared with Fenton process (Reaction (1)), UV Fenton process is becoming attractive due to the decomposition of  $\text{H}_2\text{O}_2$  and the photo-active  $\text{Fe}(\text{OH})^{2+}$  leading to additional  $\text{HO}\cdot$  radicals in solution (Javier Benitez et al., 2002; Tokumura et al., 2006), which makes Fenton process improved by the participation of photo generated  $\text{Fe}^{2+}$  and  $\text{HO}\cdot$  (Reactions (2)–(3)).



The above Fenton or photo-Fenton process for the degradation of antibiotics has been carried out at pH 3.0 (Alexy et al., 2004; Arslan-Alaton and Dogruel, 2004). However, the pH of real wastewater in many cases is neutral or alkaline, therefore pre-adjustment of pH is necessary with the added cost in chemicals and labor (Sabhi and Kiwi, 2001). Thus, it becomes an increasing concerned topic of improving the oxidation activity of Fenton or Fenton like process at ambient circumstance (Sun and Pignatello, 1992; Zhou et al., 2008).

EDTA used as catalyst for Fenton or Fenton like system to provide high efficiency for organics degradation, resulting from producing strong oxidants such as  $\text{HO}\cdot$  radicals that can quickly and nonselectively decompose organic compounds (Noradoun and Cheng, 2005; Zhou et al., 2008). A previous investigation reported by our research group demonstrated that EDTA had a higher catalytic ability for the electro- $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process, enhancing the treatment efficiency of amoxicillin (Shen et al., 2009). To date, there is no information on antibiotics treatment

\* Corresponding author. E-mail: [xmli@hnu.cn](mailto:xmli@hnu.cn)

*jesc.ac.cn*

with UV-Fe<sup>3+</sup>(EDTA)/H<sub>2</sub>O<sub>2</sub> process in either municipal or simulated antibiotics wastewater. Moreover, amoxicillin, as a semi-synthetic antibiotic, has been widely used as human and veterinary medicinal compounds for a long time, making livestock agriculture a major source of antibiotic pollution (Andreozzi et al., 2004). Therefore, it was selected as the model contamination.

This study, is expected to explore the applicability and mechanism of UV light and EDTA that contributed to the Fe<sup>3+</sup>/H<sub>2</sub>O<sub>2</sub> process. The UV-Fe<sup>3+</sup>(EDTA)/H<sub>2</sub>O<sub>2</sub> process was evaluated by the removal efficiency of amoxicillin and COD<sub>Cr</sub>, biodegradability, UV-Vis spectra and high performance liquid chromatography-mass spectra (HPLC-MS).

## 1 Materials and methods

### 1.1 Materials

H<sub>2</sub>O<sub>2</sub> (30%, W/V), 2-propanol (99%), Na<sub>2</sub>SO<sub>4</sub>, K<sub>2</sub>Ti(C<sub>2</sub>O<sub>4</sub>)<sub>3</sub>, FeCl<sub>3</sub>·7H<sub>2</sub>O, Na<sub>2</sub>EDTA (EDTA, 99.9%), H<sub>2</sub>SO<sub>4</sub>, NaOH, HCOONH<sub>4</sub> (99%), HCOOH, CH<sub>3</sub>OH, CH<sub>3</sub>CN (HPLC grade), N-Bromosuccinimide (NBS), tetrabutylammoniumbromide (TBA-Br) (AR), amoxicillin reference standard and amoxicillin capsule (C<sub>16</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>S·3H<sub>2</sub>O) were purchased from Sinopharm Chemical Reagent Co., Ltd., China. All of the reagents used without further purified. The water used in this work was deionized and purified by a pure water system (Milli-Q, Millipore, USA).

### 1.2 Experimental procedure

Reaction was conducted at 25 ± 0.2°C in a 1500-mL cylindrical Pyrex thermostatic cell having 1000 mL of working volume equipped with a magnetic stirrer. A UVA light tube (TLMINI6W, Philips) was used as light source, which was immersed into the reaction solution, providing a light intensity 6.0 mW/cm<sup>2</sup>. The experiments were performed at different pH adjusted by 1.0 mol/L H<sub>2</sub>SO<sub>4</sub> or 1.0 mol/L NaOH. Hydroxyl radical or H<sub>2</sub>O<sub>2</sub> potential reactions in the collected effluents were quenched by 6.0 mol/L NaOH (Potter and Roth, 1993).

### 1.3 Analytic methods

The samples were taken out at certain intervals from the degraded solution. The pH of solution sample was adjusted to 10.0 to remove ferric/ferrous ion and then filtered by a Millipore filter (pore size of 0.22 μm) for COD<sub>Cr</sub> measurement. The samples for UV-Visible spectra and HPLC detection were prepared by freeze-drying under vacuum at -45°C and then further dissolved in methanol and filtered by a Millipore filter (pore size of 0.22 μm).

Amoxicillin concentration was determined using a UV-Visible spectrophotometer (UV2550, Shimadzu, Japan) with a 1-cm quartz cell according to the methods presented by Saleh (1996), in which the oxidation of amoxicillin with N-bromosuccinimide (0.05%) in sodium hydroxide (0.1

mol/L) to give an intense yellow product (395 nm).

H<sub>2</sub>O<sub>2</sub> solution was prepared with 30% H<sub>2</sub>O<sub>2</sub> calibrated by KMnO<sub>4</sub> titration, which had been calibrated by Na<sub>2</sub>C<sub>2</sub>O<sub>4</sub> titration. The H<sub>2</sub>O<sub>2</sub> concentration during the reaction was quantified by UV-vis spectra at 400 nm after color was developed with potassium titanium oxalate (K<sub>2</sub>Ti(C<sub>2</sub>O<sub>4</sub>)<sub>3</sub>) in 1.0 mol/L H<sub>2</sub>SO<sub>4</sub> solution to prevent any interference caused by the organic disturbance (Sellers, 1980).

COD<sub>Cr</sub> experiments were conducted according to APHA standard methods. Biological oxygen demand (BOD<sub>5</sub>) was determined by a BOD system (OxiTop-IS6, WTW, Germany). In all biological analyses, hydrogen peroxide and iron in solution were previously removed to avoid errors in the measurement.

The concentration of Fe<sup>3+</sup>(EDTA) in sample filtrates was determined by reverse phase high performance liquid chromatography (Agilent 1100, USA) with UV light detection (λ = 258 nm, Hewlett-Packard 1050 series), which referred to the method developed by Nowack (2002) and Noradoun and Cheng (2005). The mobile phase consisted of 92% 0.02 mol/L formate buffer (formic acid sodium salt 99%, formic acid 89.5%) and 8.0% acetonitrile containing 0.001 mol/L tetrabutylammoniumbromide (TBA-Br). pH was adjusted to 3.5. The standard Fe<sup>3+</sup>(EDTA) complex was created by adding equivalent FeCl<sub>3</sub>·7H<sub>2</sub>O to the EDTA sample prior to HPLC analysis. Controls showed that neither EDTA nor Fe<sup>3+</sup> salts alone gave UV absorbance at 258 nm. All samples were filtered using 0.22 μm nylon syringe filters prior to analysis.

HPLC-MS spectra of the degraded products were carried out on LCQ-Advantage (USA) equipped with an ODS C<sub>18</sub> column (150 mm × 2.1 mm, i.d., 3.5 μm), referring to the methods reported by Nägele and Moritz (2005). The mobile phase consisting of A (water containing 10 mmol/L of ammonium formate)–B (acetonitrile containing 0.1% of formic acid) was delivered at 1.0 mL/min using a gradient elution as follows: 0–6 min of 15%–50% B; 6–17 min of 50% B; and 17–23 min of 50%–15% B. The auto-sampler was employed to make injections of 20 μL samples at room temperature. The mass spectrometers were operated under the following conditions: electrospray ionization (ESI) in positive mode with dual sprayer of ion trap MS source; 5.0 L/min of drying gas, 300°C of gas temperature; 1 kPa of nebulizer; 150,000 of ion current control; 50 msec of maximum accumulation time; 100–400 of scan; 55 L/hr of cone gas flow; 300°C of desolvation temperature; 100°C of ion source temperature; 400 L/hr of desolvation gas flow; and 3800 V of capillary.

### 1.4 HO· scavenging investigation

HO· scavenging investigation was conducted to determine whether the oxidation process was mediated by HO· in the UV-Fe<sup>3+</sup>(EDTA)/H<sub>2</sub>O<sub>2</sub> process. The molar ratio of 2-propanol (an effective scavenger of HO·) to Fe<sup>3+</sup> was selected as 50:1 (Ndjou'ou et al., 2006). Amoxicillin removal efficiency was measured at selected time intervals to evaluate the oxidation efficiency.

## 2 Results and discussion


### 2.1 Role of initial pH

As shown in Fig. 1, the highest treatment efficiency occurred at pH 3.0 in the three processes. It was evident that a change in the pH of the solution to either side of 3.0 leads to a decrease in oxidation efficiency.

In the  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process, the following mechanisms likely explained the observations: at the strong acidic medium ( $\text{pH} < 3.0$ ), the active species  $\text{Fe}(\text{H}_2\text{O})_5\text{OH}^{2+}$  was easily changed into  $\text{Fe}(\text{H}_2\text{O})_6^{2+}$ , which had lower catalysis than that of  $\text{Fe}(\text{H}_2\text{O})_5\text{OH}^{2+}$ , and thus the concentration of hydroxyl radicals  $\text{HO}\cdot$  was relative lower, decreasing the treatment efficiency (Burbano et al., 2005). When the pH raised upon 3.0, the decrease in oxidation efficiency was due to the formation of  $\text{Fe}(\text{OH})_3$ , leading to a decrease in  $\text{Fe}^{2+}$  concentration (Malik, 2004).

In the UV- $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process,  $\text{H}_2\text{O}_2$  and the photo-active  $\text{Fe}(\text{OH})^{2+}$  could lead to additional  $\text{HO}\cdot$  in solution (Reactions (2)–(3)) under the irradiation of UV light which could attack and initiate the oxidation of pollutant with relatively higher treatment efficiency than that in  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process.

In the UV- $\text{Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  process, 76.1%  $\text{TCOD}_{\text{Cr}}$  (total  $\text{COD}_{\text{Cr}}$  that amoxicillin and EDTA contribute together) and 100% amoxicillin removal efficiency could be also achieved even though the pH was increased to 7.0, while the treatment efficiency decreased to 43.0% and 59.0% in the UV- $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process, and 31.3% and 39.6% in the  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process, respectively. The results could be ascribed that EDTA and  $\text{Fe}^{3+}$  formed stable complex therefore the solubility of  $\text{Fe}^{3+}$  was improved in the aqueous solution, facilitating the production of  $\text{HO}\cdot$ ; on the other hand,  $\text{Fe}^{3+}(\text{EDTA})$  or  $\text{Fe}^{2+}(\text{EDTA})$  could react with  $\text{H}_2\text{O}_2$  quantitatively to form  $\text{HO}\cdot$  when the pH ranged from 3.0 to 8.0 (Ndjou'ou et al., 2006; Sun and Pignatello, 1992),




**Fig. 1** Initial pH effect on  $\text{COD}_{\text{Cr}}$  removal efficiency in the UV- $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  and  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  processes;  $\text{TCOD}_{\text{Cr}}$  removal efficiency in the UV- $\text{Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  process (a); amoxicillin removal efficiency in the UV- $\text{Fe}^{3+}/\text{H}_2\text{O}_2$ , UV- $\text{Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  and  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  processes (b). Experimental conditions: amoxicillin 250 mg/L,  $\text{H}_2\text{O}_2$  64.7 mmol/L,  $\text{Fe}^{3+}$  0.80 mmol/L, EDTA 0, 0.40 mmol/L, and temperature 25°C.  $\text{TCOD}_{\text{Cr}}$  means the total  $\text{COD}_{\text{Cr}}$  that amoxicillin and EDTA contribute together.

promoting the oxidation efficiency. In addition, when the pH was increased to 8.0–9.0, the UV- $\text{Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  process still exhibited higher removal efficiency than those of UV- $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  or  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  process. The result should be ascribed that the formation of  $\text{Fe}^{3+}(\text{EDTA})\text{O}_2^{3-}$ , which was a catalyst for the decomposition of hydrogen peroxide to form oxidation media, enhancing the extent of amoxicillin oxidation; whereas, when the pH raised beyond 10.5,  $\text{H}_2\text{O}_2$  was cleanly decomposed to  $\text{O}_2$  and water (Walling et al., 1970), which could not constitute an effective source of  $\text{HO}\cdot$  in aqueous solution, leading to a poor degradation of amoxicillin. Therefore, the pH for this study was selected as 7.0

### 2.2 Molar ratio of $\text{Fe}^{3+}$ to EDTA

$\text{TCOD}_{\text{Cr}}$  and amoxicillin removal efficiency were generally enhanced when the molar ratio of  $\text{Fe}^{3+}$  to EDTA increased from 1:4 to 2:1 (Fig. 2). The results indicated that the molar ratio of  $\text{Fe}^{3+}$  to EDTA was responsible for the formation of  $\text{Fe}^{3+}$ -EDTA complex, which played a crucial role in the equilibrium among the oxidation, reduction and coordination reactions (Noradoun and Cheng, 2005). The excess EDTA would produce an extra contamination and also act as an antioxidant (Nicoli et al., 2000) that enhanced the activity of phenolic hydroxyl of amoxicillin, improving its anti-oxidation ability. Whereas, the treatment efficiency decreased when the molar ratio of  $\text{Fe}^{3+}$  to EDTA changed from 2:1 to 4:1 because the over dosage of  $\text{Fe}^{3+}$  acted as scavenger of  $\text{H}_2\text{O}_2$  to form less reactive  $\text{HO}_2\cdot$ . Another possible explanation was that the excess of  $\text{Fe}^{3+}$  would lead to the formation of  $\text{Fe}(\text{OH})_3$ , destroying the complex and reducing the generation rate of  $\text{HO}\cdot$ . Therefore, the optimal molar ratio of  $\text{Fe}^{3+}$  to EDTA was selected as 2:1. It should be noted that the dosage of EDTA was halved in the UV- $\text{Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  process, compared with that in the previous investigation (Shen et al., 2009). The results should be attributed that  $\text{Fe}^{3+}$  or  $\text{Fe}(\text{OH})^{2+}$  was constantly photo-generated  $\text{Fe}^{2+}$  under irradiation of UV light (Reactions (3)–(4)) and inhibited the formation of  $\text{Fe}(\text{OH})_3$ , resulting in the decrease of the



**Fig. 2**  $\text{TCOD}_{\text{Cr}}$  (a) and amoxicillin (b) removal efficiency under varying molar ratios of  $\text{Fe}^{3+}$  to EDTA in the UV- $\text{Fe}^{3+}(\text{EDTA})/\text{H}_2\text{O}_2$  process. Experimental conditions: amoxicillin 250 mg/L,  $\text{H}_2\text{O}_2$  64.7 mmol/L,  $\text{Fe}^{3+}$  0.80 mmol/L, EDTA<sub>0</sub> 0.18 mol/L, pH 7.0, and temperature 25°C.


EDTA dosage. Therefore, the optimum molar ratio of  $\text{Fe}^{3+}$  to EDTA was selected as 2:1, and the concentration of EDTA was 0.40 mmol/L.

### 2.3 Molar ratio of $\text{H}_2\text{O}_2$ to $\text{Fe}^{3+}$


Figure 3 shows that  $\text{TCOD}_{\text{Cr}}$  and amoxicillin removal efficiency increased when the molar ratio of  $\text{H}_2\text{O}_2$  to  $\text{Fe}^{3+}$  increased from 36.0 to 108. However, when the molar ratio of  $\text{H}_2\text{O}_2$  to  $\text{Fe}^{3+}$  was larger than 80.9, it could not lead to significant increase in the degradation efficiency. This might be attributed that the unconsumed  $\text{H}_2\text{O}_2$  served as a scavenger for  $\text{HO}\cdot$  and resulted in less reactive  $\text{HO}_2\cdot$  (Zhou et al. 2007), decreasing amoxicillin degradation. Additionally, it should be noted that the residual  $\text{H}_2\text{O}_2$  could create extra  $\text{COD}_{\text{Cr}}$ . Therefore, the optimum molar ratio of  $\text{H}_2\text{O}_2$  to  $\text{Fe}^{3+}$  was selected as 80.9, and the concentration of  $\text{H}_2\text{O}_2$  was 64.7 mmol/L.

### 2.4 $\text{COD}_{\text{Cr}}$ distribution between EDTA and amoxicillin

As shown in Fig. 4, EDTA was degraded in the oxidation process due to the decreasing concentration of  $\text{Fe}^{3+}$ (EDTA) and its  $\text{COD}_{\text{Cr}}$  removal efficiency. It was



**Fig. 3**  $\text{TCOD}_{\text{Cr}}$  (a) and amoxicillin (b) removal efficiency under varying molar ratios of  $\text{H}_2\text{O}_2$  to  $\text{Fe}^{3+}$ . Experimental conditions: amoxicillin 250 mg/L,  $\text{H}_2\text{O}_2^0$  32.3 mol/L,  $\text{Fe}^{3+}$  0.80 mmol/L, EDTA 0.40 mmol/L, pH 7.0, and temperature 25°C.








**Fig. 4** Variation of  $\text{Fe}^{3+}$ (EDTA) concentration (a) and  $\text{COD}_{\text{Cr}}$  removal efficiency of EDTA (b) in the system that EDTA was selected as the sole contamination. Experimental conditions:  $\text{Fe}^{3+}$  0.80 mmol/L, EDTA 0.40 mmol/L,  $\text{H}_2\text{O}_2$  64.7 mmol/L, pH 7.0, and temperature 25°C.

therefore that both EDTA and amoxicillin contributed to  $\text{COD}_{\text{Cr}}$  removal efficiency ( $\text{TCOD}_{\text{Cr}}$ ), yet which was difficult to be distinguished quantitatively only by the  $\text{UV-Fe}^{3+}$ (EDTA)/ $\text{H}_2\text{O}_2$  process. Therefore, a control experiment that EDTA selected as the sole contamination was conducted under the same experimental conditions. The removal efficiency of  $\text{TCOD}_{\text{Cr}}$  was found to be 76.1% at pH 7.0 and the initial  $\text{COD}_{\text{Cr}}$  of amoxicillin (0.25 g/L) and EDTA (0.40 mmol/L) were 360.0 and 135.0 mg/L, respectively. The total  $\text{COD}_{\text{Cr}}$  removal could be calculated as 376.7 mg/L according to the total  $\text{COD}_{\text{Cr}}$  multiplying by the total removal efficiency. Similarly, 81.7 mg/L of  $\text{COD}_{\text{Cr}}$  removal of EDTA could be obtained referring to 60.5% removal efficiency in Fig. 4b. Therefore, the  $\text{COD}_{\text{Cr}}$  removal of amoxicillin and the  $\text{COD}_{\text{Cr}}$  removal efficiency could be calculated as 295 mg/L and 81.9%, respectively.

### 2.5 Mechanisms investigation


#### 2.5.1 Catalytic mechanisms of EDTA and UV light

In the  $\text{UV-Fe}^{3+}$ (EDTA)/ $\text{H}_2\text{O}_2$  process, EDTA was employed to enhance iron cycles that catalyzed  $\text{H}_2\text{O}_2$  to produce  $\text{HO}\cdot$  or other activated media (Sun and Pignatello, 1992; Zhou et al., 2008). The efficiency of  $\text{HO}\cdot$  formation increased by the photo-reaction of  $\text{H}_2\text{O}_2$  and/or  $\text{Fe}(\text{OH})^{2+}$  that produced  $\text{HO}\cdot$  directly or regenerated  $\text{Fe}^{2+}$  to feed back the chain reaction (Reactions (1)–(6)) (Javier Benitez et al., 2002; Tokumura et al., 2006). UV acceleration was most likely due to charge-transfer photoreduction of  $\text{Fe}^{3+}$ (EDTA) to give  $\text{Fe}^{2+}$ , which reacted with  $\text{H}_2\text{O}_2$  to generate hydroxyl radical in the Fenton reaction (Sun and Pignatello, 1993). On the other hand, the enhancement of treatment efficiency might be ascribed to a photo-induced oxidation by an inner-sphere electron-transfer reaction of the ligands of various  $\text{Fe}(\text{III})$  complexes such as  $\text{Fe}^{3+}$ (EDTA),  $\text{Fe}(\text{OH})_3$  with  $\text{H}_2\text{O}$ , the organic substrate, or its intermediates of degradation (Bossman et al., 1998). Therefore, the following mechanisms could be further derived from the investigation.



As shown in Fig. 5a,  $\text{H}_2\text{O}_2$  concentration gradually decreased with the on-going oxidation process. It could be accounted that the large amount of  $\text{H}_2\text{O}_2$  contributed to producing  $\text{HO}\cdot/\text{HO}_2\cdot$  under the irradiation of UV light or reacting with  $\text{Fe}^{3+}$ (EDTA)/ $\text{Fe}^{2+}$ (EDTA) (Eqs. (2), (4) and (5)).

Figure 5b presents that the pH decreased gradually from 7.0 to 3.0 in the oxidation process, which implied that low molecule weight organic acids or acidic media might be produced during the oxidation process. On the other hand, the decreasing pH was helpful for EDTA degradation because EDTA was subject to Fenton oxidation at acidic conditions (Walling et al., 1970). It was manifested that the initial higher pH inhibited the oxidation of EDTA during



**Fig. 5**  $\text{H}_2\text{O}_2$  decay rate (a); pH changes (b); and amoxicillin removal efficiency (c) in the UV- $\text{Fe}^{3+}$ (EDTA)/ $\text{H}_2\text{O}_2$  process and UV- $\text{Fe}^{3+}$ (EDTA)/ $\text{H}_2\text{O}_2$  process scavenged by 2-propanol. Experimental conditions: amoxicillin 250 mg/L,  $\text{H}_2\text{O}_2$  64.7 mol/L,  $\text{Fe}^{3+}$  0.80 mmol/L, EDTA 0.40 mmol/L, pH 7.0, and temperature 25°C.


the initial reaction step, even though there was a large amount of  $\text{HO}\cdot$ . Therefore, EDTA first acted as catalyst and then as reactant when the pH decreased to 3.0.

Figure 5c showed that the degradation of amoxicillin was dramatically decreased to 18.7% when 2-propanol ( $\text{HO}\cdot$  scavenger) was added into the UV- $\text{Fe}^{3+}$ (EDTA)/ $\text{H}_2\text{O}_2$  process. These results demonstrated that  $\text{HO}\cdot$  was a primary reactant in the oxidation process. It was also revealed that other activated media might exist in the oxidation process, leading to the tiny degradation of amoxicillin.

### 2.5.2 Potential activated media in the UV- $\text{Fe}^{3+}$ (EDTA)/ $\text{H}_2\text{O}_2$ process

The multi-staged degradation curves (Figs. 2, 3) proposed that EDTA,  $\text{H}_2\text{O}_2$ , and  $\text{Fe}^{3+}$  might build new activated media, which could be responsible for enhancing  $\text{COD}_{\text{Cr}}$  and amoxicillin removal, promoting the oxidation process. To confirm the resulted activated media, UV-Vis spectra of the following solutions were performed. As shown in Fig. 6a, only EDTA didn't exhibit any specific absorption at the range of 200–800 nm, and  $\text{Fe}^{3+}$  exhibited characteristic absorption at 223 and 301.5 nm, whereas  $\text{Fe}^{3+}$ (EDTA) led to a new maximum absorption at 258 nm. Figure 6b further shows that the mixture of  $\text{Fe}^{3+}$  and  $\text{H}_2\text{O}_2$  led to a new maximum absorption at 303 nm, and alone  $\text{H}_2\text{O}_2$  without any specific absorption. Amoxicillin caused maximum absorptions at 230 and 272 nm, whereas the mixture of  $\text{Fe}^{3+}$  and amoxicillin exhibited maximum absorptions at 229.5 and 278 nm. Therefore, these new absorptions likely contributed to the formation of new activated media, which cooperated with  $\text{HO}\cdot$  for the degradation of amoxicillin.


The results of  $\text{BOD}_5/\text{COD}_{\text{Cr}}$  further revealed that the UV- $\text{Fe}^{3+}$ (EDTA)/ $\text{H}_2\text{O}_2$  process was a promising strategy to degrade amoxicillin as the biodegradability of the effluent was improved to 0.45, compared with the cases of UV- $\text{Fe}^{3+}/\text{H}_2\text{O}_2$ (0.25) and  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  processes (0.10) (Fig. 6c).



**Fig. 6** UV-Vis spectra of  $\text{Fe}^{3+}$ , EDTA, initial  $\text{Fe}^{3+}$ (EDTA), and degraded  $\text{Fe}^{3+}$ (EDTA) (a);  $\text{Fe}^{3+}$ , amoxicillin,  $\text{H}_2\text{O}_2$ , the mixture of  $\text{Fe}^{3+}$  and amoxicillin,  $\text{Fe}^{3+}$  and  $\text{H}_2\text{O}_2$  (b);  $\text{BOD}_5/\text{COD}_{\text{Cr}}$  in the UV- $\text{Fe}^{3+}$ (EDTA)/ $\text{H}_2\text{O}_2$  (I), UV- $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  (II) and  $\text{Fe}^{3+}/\text{H}_2\text{O}_2$  (III) processes (the insert plot in Fig. 6b). Experimental conditions: amoxicillin 250 mg/L,  $\text{H}_2\text{O}_2$  64.7 mmol/L,  $\text{Fe}^{3+}$  0.80 mmol/L, EDTA 0.40 mmol/L, pH 7.0, and temperature 25°C.

### 2.5.3 Degradation pathway of amoxicillin

The degraded products of amoxicillin were separated by HPLC at the retention time of 1.53, 2.63, and 16.13 min (Fig. 7), and each peak was characterized by its mass measurements ( $m/z$ ), respectively (Fig. 8). The degradation of amoxicillin started with the opening of the four-membered  $\beta$ -lactam ring and yielded the product amoxicillin penicilloic acid at  $m/z$  383 ( $\text{C}_{16}\text{H}_{22}\text{N}_3\text{O}_6\text{S}$ ) (Figs. 8b, 9a) which contained a free carboxylic acid group providing a higher polarity to this molecule (Nägele and Moritz, 2005). The structure of this degradation product was confirmed by the appearance of the characteristic fragment at  $m/z$  340 ( $\text{C}_{15}\text{H}_{22}\text{N}_3\text{O}_4\text{S}$ ) (Fig. 8b), which is the product of a decarboxylation reaction and the fragment at  $m/z$  189 ( $\text{C}_7\text{H}_{13}\text{N}_2\text{O}_2\text{S}$ ) was another important proof for the structure confirmation (Fig. 8b). Following by amoxicillin penicilloic acid, there were two possible pathways for the further degradation. The first one was based on the decarboxylation of the free carboxylic acid and demethyl



**Fig. 7** HPLC chromatogram of the degraded amoxicillin.




Fig. 8 Mass spectra of the degraded amoxicillin. Experimental conditions: ESI in positive model with scan of  $m/z$  100–400.




Fig. 9 Degradation pathway of amoxicillin.

reaction of the five-membered thiazolidine ring leading to the compound amoxicillin penicilloic acid at  $m/z$  325 ( $C_{14}H_{19}N_3O_4S$ ) (Figs. 8a, 9b). The second one was undergone cleavage of the five-membered thiazolidine ring and decarboxylation yielding amoxicillin penicillamine at  $m/z$  284 ( $C_{12}H_{16}N_2O_4S$ ) (Figs. 8c, 9c). It was possible that another intermediate products was the formation of a new, stable, six-membered ring giving diketopiperazine amoxicillin (the isomeric compound of amoxicillin) at  $m/z$  365 ( $C_{16}H_{19}N_3O_5S$ ) (Figs. 8b, 9d).

### 3 Conclusions

In this work, a novel oxidation process of UV- $Fe^{3+}$ (EDTA)/ $H_2O_2$  has been investigated for the degradation of amoxicillin at pH 7.0, resulting in appreciable treatment efficiency. It was found that EDTA and UV light performed catalytic effect on the oxidation process synergistically, which involved in multi-step mechanisms of Fenton, Fenton like, photo-Fenton and Fe(III)-organic complexes. The results further revealed that EDTA acted

as catalyst and reactant as well.

## Acknowledgments

The work was supported by the National Science Foundation of China (No. 51078128) and the Hunan Provincial Innovation Foundation for Postgraduate (No. CX2009B079).

## References

- Alexy R, Kumpel T, Kümmerer K, 2004. Assessment of degradation of 18 antibiotics in the closed bottle test. *Chemosphere*, 57(6): 505–512.
- Andreozzi R, Caprio V, Ciniglia C, de Champdoré M, Giudice R L, Marotta R et al., 2004. Antibiotics in the environment: occurrence in Italian STPs, fate, and preliminary assessment on algal toxicity of amoxicillin. *Environmental Science and Technology*, 38(24): 6832–6838.
- Arslan-Alaton I, Dogruel S, 2004. Pre-treatment of penicillin formulation effluent by advanced oxidation processes. *Journal of Hazardous Materials*, 112(12): 105–113.
- Bossmann S H, Oliveros E, Göb S, Siegwart S, Dahlen E P, Payawan L Jr et al., 1998. New evidence against hydroxyl radicals as reactive intermediates in the thermal and photochemically enhanced Fenton reactions. *The Journal of Physical Chemistry A*, 102(28): 5542–5550.
- Burbano A A, Dionysiou D D, Suidan M T, Richardson T L, 2005. Oxidation kinetics and effect of pH on the degradation of MTBE with Fenton reagent. *Water Research*, 39(1): 107–118.
- da Silva M F, Tiago I, Veríssimo A, Boaventura R A R, Nunes O C, Manaia C M, 2006. Antibiotic resistance of enterococci and related bacteria in an urban wastewater treatment plant. *FEMS Microbiology Ecology*, 55(2): 322–329.
- Gulkowska A, Leung H W, So M K, Taniyasu S, Yamashita N, Yeung L W Y et al., 2008. Removal of antibiotics from wastewater by sewage treatment facilities in Hong Kong and Shenzhen, China. *Water Research*, 42(1-2): 395–403.
- Huber M M, Canonica S, Park G Y, von Gunten U, 2003. Oxidation of pharmaceuticals during ozonation and advanced oxidation processes. *Environmental Science and Technology*, 37(5): 1016–1024.
- Javier Benitez F, Acero J L, Real F J, 2002. Degradation of carbofuran by using ozone, UV radiation and advanced oxidation processes. *Journal of Hazardous Materials*, 89(1): 51–65.
- Kümmerer K, Al-Ahmad A, Mersch-Sundermann V, 2000. Biodegradability of some antibiotics, elimination of the genotoxicity and affection of wastewater bacteria in a simple test. *Chemosphere*, 40(7): 701–710.
- Malik P K, 2004. Oxidation of safranin T in aqueous solution using Fenton's reagent: involvement of an Fe(III) chelate in the catalytic hydrogen peroxide oxidation of safranin T. *The Journal of Physical Chemistry A*, 108(14): 2675–2681.
- Migliore L, Civitareale C, Brambilla G, Di Delupis G D, 1997. Toxicity of several important agricultural antibiotics to *Artemia*. *Water Research*, 31(7): 1801–1806.
- Nägele E, Moritz R, 2005. Structure elucidation of degradation products of the antibiotic amoxicillin with ion trap MS<sup>n</sup> and accurate mass determination by ESI TOF. *Journal of the American Society for Mass Spectrometry*, 16(10): 1670–1676.
- Ndjou'ou A C, Bou-Nasr J, Cassidy D, 2006. Effect of Fenton reagent dose on coexisting chemical and microbial oxidation in soil. *Environmental Science and Technology*, 40(8): 2778–2783.
- Nicoli M C, Calligaris S, Manzocco L, 2000. Effect of enzymatic and chemical oxidation on the antioxidant capacity of catechin model systems and apple derivatives. *Journal of Agricultural and Food Chemistry*, 48(10): 4576–4580.
- Noradoun C E, Cheng I F, 2005. EDTA degradation induced by oxygen activation in a zerovalent iron/air/water system. *Environmental Science and Technology*, 39(18): 7158–7163.
- Nowack B, 2002. Environmental chemistry of aminopolycarboxylate chelating agents. *Environmental Science and Technology*, 36(19): 4009–4016.
- Potter F J, Roth J A, 1993. Oxidation of chlorinated phenols using Fenton's reagent. *Hazardous Waste and Hazardous Materials*, 10(2): 151–170.
- Sabhi S, Kiwi J, 2001. Degradation of 2,4-dichlorophenol by immobilized iron catalysts. *Water Research*, 35(8): 1994–2002.
- Saleh G A, 1996. Two selective spectrophotometric methods for the determination of amoxicillin and cefadroxil. *Analyst*, 121(5): 641–645.
- Sellers R M, 1980. Spectrophotometric determination of hydrogen peroxide using potassium titanium(IV) oxalate. *Analyst*, 105(1255): 950–954.
- Shen T T, Li X M, Tang Y F, Wang J, Yue X, Cao J B et al., 2009. EDTA and electricity synergetic catalyzed Fe<sup>3+</sup>/H<sub>2</sub>O<sub>2</sub> process for amoxicillin oxidation. *Water Science and Technology*, 60(3): 761–770.
- Sun Y F, Pignatello J J, 1992. Chemical treatment of pesticide wastes: Evaluation of iron(III) chelates for catalytic hydrogen peroxide oxidation of 2,4-D at circumneutral pH. *Journal of Agricultural and Food Chemistry*, 40(2): 322–327.
- Sun Y F, Pignatello J J, 1993. Activation of hydrogen peroxide by iron(III) chelates for abiotic degradation of herbicides and insecticides in water. *Journal of Agricultural and Food Chemistry*, 41(2): 308–312.
- Tokumura M, Ohta A, Znad H T, Kawase Y, 2006. UV light assisted decolorization of dark brown colored coffee effluent by photo-Fenton reaction. *Water Research*, 40(20): 3775–3784.
- Walling C, Kurz M, Schugar H J, 1970. Iron(III)-ethylenediaminetetraacetic acid-peroxide system. *Inorganic Chemistry*, 9(4): 931–937.
- Zhou M H, Yu Q H, Lei L C, Barton G, 2007. Electro-Fenton method for the removal of methyl red in an efficient electrochemical system. *Separation and Purification Technology*, 57(2): 380–387.
- Zhou T, Li Y Z, Wong F S, Lu X H, 2008. Enhanced degradation of 2,4-dichlorophenol by ultrasound in a new Fenton like system (Fe/EDTA) at ambient circumstance. *Ultrasonics Sonochemistry*, 15(5): 782–790.