


JES

JOURNAL OF
ENVIRONMENTAL
SCIENCES

ISSN 1001-0742
CN 11-2629/X

December 1, 2014 Volume 26 Number 12
www.jesc.ac.cn



Sponsored by
Research Center for Eco-Environmental Sciences
Chinese Academy of Sciences

- 2369 Effects of seasonal climatic variability on several toxic contaminants in urban lakes: Implications for the impacts of climate change
Qiong Wu, Xinghui Xia, Xinli Mou, Baotong Zhu, Pujun Zhao, and Haiyang Dong
- 2379 Preparation of cross-linked magnetic chitosan with quaternary ammonium and its application for Cr(VI) and P(V) removal
Wei Yao, Pinhua Rao, Irene M.C. Lo, Wenqi Zhang, and Wenrui Zheng
- 2387 Formation pathways of brominated products from benzophenone-4 chlorination in the presence of bromide ions
Ming Xiao, Dongbin Wei, Liping Li, Qi Liu, Huimin Zhao, and Yuguo Du
- 2397 Influence of the inherent properties of drinking water treatment residuals on their phosphorus adsorption capacities
Leilei Bai, Changhui Wang, Liansheng He, and Yuansheng Pei
- 2406 Radiation induced decomposition of a refractory cefthiamidine intermediate
Qiburi Bao, Lujun Chen, and Jianlong Wang
- 2412 Characterization of aerosol optical properties, chemical composition and mixing states in the winter season in Shanghai, China
Yong Tang, Yuanlong Huang, Ling Li, Hong Chen, Jianmin Chen, Xin Yang, Song Gao, and Deborah S. Gross
- 2423 Knudsen cell and smog chamber study of the heterogeneous uptake of sulfur dioxide on Chinese mineral dust
Li Zhou, Weigang Wang, Yanbo Gai, and Maofa Ge
- 2434 Experimental study on filtration and continuous regeneration of a particulate filter system for heavy-duty diesel engines
Tao Tang, Jun Zhang, Dongxiao Cao, Shijin Shuai, and Yanguang Zhao
- 2440 Combination of heterogeneous Fenton-like reaction and photocatalysis using Co-TiO₂ nanocatalyst for activation of KHSO₅ with visible light irradiation at ambient conditions
Qingkong Chen, Fangying Ji, Qian Guo, Jianping Fan, and Xuan Xu
- 2451 Atmospheric sulfur hexafluoride *in-situ* measurements at the Shangdianzi regional background station in China
Bo Yao, Lingxi Zhou, Lingjun Xia, Gen Zhang, Lifeng Guo, Zhao Liu, and Shuangxi Fang
- 2459 Direct radiative forcing of urban aerosols over Pretoria (25.75°S, 28.28°E) using AERONET Sunphotometer data: First scientific results and environmental impact
Ayodele Joseph Adesina, Kanike Raghavendra Kumar, Venkataraman Sivakumar, and Derek Griffith
- 2475 Chemical characteristics and source apportionment of atmospheric particles during heating period in Harbin, China
Likun Huang and Guangzhi Wang
- 2484 Microbial community structures in an integrated two-phase anaerobic bioreactor fed by fruit vegetable wastes and wheat straw
Chong Wang, Jiane Zuo, Xiaojie Chen, Wei Xing, Linan Xing, Peng Li, Xiangyang Lu, and Chao Li
- 2493 Persistent pollutants and the patchiness of urban green areas as drivers of genetic richness in the epiphytic moss *Leptodon smithii*
Valeria Spagnuolo, Flavia De Nicola, Stefano Terracciano, Roberto Bargagli, Daniela Baldantoni, Fabrizio Monaci, Anna Alfani, and Simonetta Giordano

CONTENTS

- 2500 Enhanced removal of ethylbenzene from gas streams in biotrickling filters by Tween-20 and Zn(II)
Lu Wang, Chunping Yang, Yan Cheng, Jian Huang, Haining Yang, Guangming Zeng, Li Lu, and Shanying He
- 2508 Enhanced efficiency of cadmium removal by *Boehmeria nivea* (L.) Gaud. in the presence of exogenous citric and oxalic acids
Huaying Li, Yunguo Liu, Guangming Zeng, Lu Zhou, Xin Wang, Yaqin Wang, Chunlin Wang, Xinjiang Hu, and Weihua Xu
- 2517 Comparative sorption and desorption behaviors of PFHxS and PFOS on sequentially extracted humic substances
Lixia Zhao, Yifeng Zhang, Shuhong Fang, Lingyan Zhu, and Zhengtao Liu
- 2526 Inhibitory effects of nisin-coated multi-walled carbon nanotube sheet on biofilm formation from *Bacillus anthracis* spores
Xiuli Dong, Eric McCoy, Mei Zhang, and Liju Yang
- 2535 A comparative study and evaluation of sulfamethoxazole adsorption onto organo-montmorillonites
Laifu Lu, Manglai Gao, Zheng Gu, Senfeng Yang, and Yuening Liu
- 2546 Removal of formaldehyde over $Mn_xCe_{1-x}O_2$ catalysts: Thermal catalytic oxidation *versus* ozone catalytic oxidation
Jia Wei Li, Kuan Lun Pan, Sheng Jen Yu, Shaw Yi Yan, and Moo Been Chang
- 2554 Humic acid transport in saturated porous media: Influence of flow velocity and influent concentration
Xiaorong Wei, Mingan Shao, Lina Du, and Robert Horton
- 2562 Salinity influence on soil microbial respiration rate of wetland in the Yangtze River estuary through changing microbial community
Xue Fei Xi, Lei Wang, Jia Jun Hu, Yu Shu Tang, Yu Hu, Xiao Hua Fu, Ying Sun, Yiu Fai Tsang, Yan Nan Zhang, and Jin Hai Chen
- 2571 Comments on “Adsorption of 2-mercaptobenzothiazole from aqueous solution by organo-bentonite” by P. Jing, M.H. Hou, P. Zhao, X.Y. Tang, H.F. Wan
Yuhshan Ho
- 2573 Reply to comments on “Adsorption of 2-mercaptobenzothiazole from aqueous solution by organo-bentonite” by Yuhshan Ho
Ping Jing, Meifang Hou, Ping Zhao, Xiaoyan Tang, and Hongfu Wan

Available online at www.sciencedirect.com

ScienceDirect

www.journals.elsevier.com/journal-of-environmental-sciences

Enhanced efficiency of cadmium removal by *Boehmeria nivea* (L.) Gaud. in the presence of exogenous citric and oxalic acids

Huaying Li^{1,2}, Yunguo Liu^{1,2,*}, Guangming Zeng^{1,2}, Lu Zhou^{1,2}, Xin Wang³, Yaqin Wang^{1,2}, Chunlin Wang^{1,2}, Xinjiang Hu^{1,2}, Weihua Xu^{1,2}

1. College of Environmental Science and Engineering, Hunan University, Changsha 410082, China. E-mail: lihuaying@163.com

2. Key Laboratory of Environmental Biology and Pollution Control, Hunan University, Ministry of Education, Changsha 410082, China

3. College of Resources and Environmental Science, Hunan Normal University, Changsha 410081, China

ARTICLE INFO

Article history:

Received 6 January 2014

Revised 14 April 2014

Accepted 11 May 2014

Available online 1 November 2014

Keywords:

Heavy metal

Phytoremediation

Lipid peroxidation

Antioxidant defense

Translocation factor

ABSTRACT

Boehmeria nivea (L.) Gaud. is a potential candidate for the remediation of Cd contaminated sites. The present investigation aims to explore Cd tolerance threshold and to quickly identify the role of exogenous organic acids in Cd uptake and abiotic metal stress damage. Elevated Cd levels (0–10 mg/L) resulted in an obvious rise in Cd accumulation, ranging from 268.0 to 374.4 in root and 25.2 to 41.2 mg/kg dry weight in shoot, respectively. Citric acid at 1.5 mmol/L significantly facilitated Cd uptake by 26.7% in root and by 1-fold in shoot, respectively. Cd translocation efficiency from root to shoot was improved by a maximum of 66.4% under 3 mmol/L of oxalic acid. Citric acid exhibited more prominent mitigating effect than oxalic acid due to its stronger ligand affinity for chelating with metal and avoiding the toxicity injury of free Cd ions more efficiently. The present work provides a potential strategy for efficient Cd remediation with *B. nivea*.

© 2014 The Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences.

Published by Elsevier B.V.

Introduction

Industrial mining, smelting and agricultural activities such as long-term wastewater irrigation have caused prevalent soil Cd contamination and hazardous human diseases through edible plants. The persistence of Cd in soils and transfer into human body via food chain have aroused a major concern due to its great toxicity (Liu et al., 2008), mutagenicity, carcinogenicity and high risk to food safety. Cd contamination is extremely severe in Hunan Province (in central-south of China) because of frequent mining activities, which has led to farmland pollution, reduction of grain production and economic loss. For example, 36% of rice grown in Hunan Province was detected to have Cd levels above China's food standard regulation according to a research report (Lei et al., 2010). In order to alleviate the problem, efficient phytoremediation seems to be an appropriate route. *Boehmeria nivea* (L.) Gaud. (ramie), a perennial Urticaceae species and a

primary fiber crop for textile as well as a dominant plant of mining sites in Hunan Province, has been identified as a new Cd tolerance plant species with deep root and high biomass (Wang et al., 2008). Compared to other herbaceous and woody plants, ramie has great superiority in phytoremediation of Cd contaminated sites especially the farmland for its particular capability of Cd accumulation, fast growth, ease of cultivation and propagation. All the above makes ramie not only an advantageous plant species to achieve economic and eco-environmental benefits, but also a valuable organism for studying Cd-induced physiological mechanisms to strengthen plant abiotic stress tolerance.

Due to its strong phytotoxicity, Cd can pose adverse symptoms on plants such as leaf chlorosis, browning of root tips and growth retardation, even resulting in generation of reactive oxygen species (ROS), inhibition of photosynthesis (López-Millán et al., 2009), membrane lipid peroxidation (Li et al., 2012), protein degradation, ultra-structural damage and ultimately cell death

* Corresponding author. E-mail: yunguoliu_hnu@163.com (Yunguo Liu).

(Daud et al., 2009). To survive Cd-induced stress, plants have evolved protective mechanisms to mitigate and repair the oxidative impairment (Edreva, 2005), including morphological changes, physiological adaptations and antioxidative defense system. Generally, the antioxidative defense mechanisms include antioxidative enzymes such as superoxide dismutase (SOD), peroxidase (POD), catalase (CAT), ascorbate peroxidase (APX) and glutathione reductase (GR), and non-enzymatic antioxidants like glutathione (GSH), ascorbic acid (AsA) and carotenoids (Fernández et al., 2013). SOD is the key enzyme involved in catalyzing the dismutation of highly reactive superoxide radicals $O_2^{\cdot -}$ to O_2 and H_2O_2 , which is further decomposed to non-toxic forms like O_2 and H_2O by APX of the ascorbate–glutathione cycle or by GPX and CAT in the cytoplasm and other cellular compartments (Jin et al., 2008).

Phytoextraction is defined as the removal of contaminants from soil by plant roots and their subsequent translocation to harvestable shoot tissues (Clabeaux et al., 2011). As an eco-friendly and cost-effective remediation technology, using plants to extract toxic metals from soil can be less destructive to soil structure and plant rhizosphere, more esthetic and more suited to impacted communities than conventional methods (Meighan et al., 2011). The extraction efficiency was closely related with contaminant availability for root uptake and translocation to the aerial parts (Bhargava et al., 2012). However, low bioavailability and limited translocation of some metals become the predominant restrictions in phytoextraction (Najeeb et al., 2011; Yang et al., 2013). Then high biomass yielding plant species with chemically assisted phytoextraction such as adding chelating agents have been used to enhance metal extraction efficiency. Chelators like low-molecular-weight organic acids (LMWOA) are capable of forming chemical complexes with metal ions and modifying the bioavailability of heavy metals in soil. LMWOA (such as citric, oxalic and malic acid), natural compounds derived from root exudates, are characterized as lower toxicity and higher biodegradability, which makes them more suitable for assisted phytoremediation than synthetic chelators.

Extensive studies have indicated that organic acids are involved in tolerance, transport and storage of heavy metal and played a key role in maintaining cellular homeostasis (Irtelli and Navari-Izzo, 2006). In particular, 25 $\mu\text{mol/L}$ of citric acid could enhance Cd uptake and translocation by 61% and 2.2-fold at 4 $\mu\text{mol/L}$ Cd level in *Halimione portulacoides*, respectively (Duarte et al., 2007). It could also counter Cd toxicity by improving plant growth, restoring shape and structure of root cells, and eliminating plasmolysis in *Juncus effusus* L. (Najeeb et al., 2011). Najeeb et al. (2009) found that citric acid was capable of enhancing Mn phytoextraction by 20% and alleviating metal toxicity from the reduced number of plastoglobuli in plant chloroplast. Similarly, Jean et al. (2008) reported that citric acid strengthened Cr and Ni uptake and translocation in *Datura innoxia*. Citric acid triggered more positive impact on Cu bioavailability and a visibly higher concentration of Cu uptake in tobacco shoot compared to oxalic and tartaric acid (Evangelou et al., 2006). Comparatively, citric acid (10 mmol/L) demonstrated a higher Cu mobilization (from 1 in the control to 42 mg/kg) than tartaric acid (Pérez-Esteban et al., 2013). These studies put forward an insight into the positive effect of exogenous organic acids on plant biomass, metal uptake and translocation, and metal tolerance for improving phytoremediation efficiency. In this work, a new Cd tolerant plant was studied to extend the range of plant species for better understanding of its potential application in phytoremediation.

Previous studies mainly focused on the accumulation, transport and tolerance mechanisms of ramie and subcellular distribution of heavy metals in plant tissues (Liu et al., 2007; Xia et al., 2009; Wang et al., 2008, 2011). However, to our best knowledge, little information is available on the effect of LMWOA on Cd phytoextraction and tolerance. Therefore it seems significant to demonstrate the natural chelator–Cd–plant interaction in this species. The objectives of this study were to: (1) explore the potential of ramie in phytoextraction of Cd; (2) compare the effect of different concentrations of exogenous citric and oxalic acids on plant growth, Cd uptake and

translocation; and (3) determine the role of citric and oxalic acids in Cd tolerance of ramie by regulating the antioxidant defense system involved in stress endurance.

1. Materials and methods

1.1. Plant collection and hydroponic culture

A batch of ramie seedlings was collected from Ramie Research Institute (Changsha, Hunan Province, China). Plants were acclimatized to hydroponic condition in 1/8 Hoagland nutrient solution ($\text{pH } 6.0 \pm 0.5$) for 2 weeks. The solution was aerated continuously and renewed every 2 days. The experiments were conducted in a growth chamber with 25/20°C, day/night temperature, 60%–70% relative humidity and 14 hr photoperiod at light intensity of 300 $\mu\text{mol}/(\text{m}^2 \cdot \text{sec})$.

1.2. Cadmium treatment and the addition of organic acids

After preculturing for 2 weeks, the plants were treated in triplicates in a completely randomized design as follows: 0 (Cd_0), 2 (Cd_2), 5 (Cd_5), 10 (Cd_{10}) mg/L Cd concentrations as $\text{Cd}(\text{NO}_3)_2$ and $\text{Cd}_{10} + \text{CA}_1$, $\text{Cd}_{10} + \text{CA}_2$, $\text{Cd}_{10} + \text{OA}_1$, $\text{Cd}_{10} + \text{OA}_2$ (CA_1 and CA_2 refer to 1.5, 4 mmol/L citric acid; OA_1 and OA_2 refer to 3, 9 mmol/L oxalic acid, respectively). Organic acids were applied at highest Cd level (10 mg/L) after Cd exposure for 1 week. Plant tissues were harvested after one month. Fresh samples were frozen immediately in liquid nitrogen and stored at -80°C for subsequent analysis.

1.3. Analysis of cadmium uptake

Plant roots and shoots were dried at 80°C till constant weight. About 0.5 g dry samples were digested with 10 mL HNO_3 and 3 mL HClO_4 at 160°C by heating to be transparent. Digested liquid was washed several times into a 100 mL volumetric flask and diluted with de-ionized water to the calibration line for measurement. The Cd content was determined by atomic absorption spectrometer (Analyst 700, PerkinElmer, Massachusetts, USA) and calculated as mg/kg of dry weight (dw). Translocation factor (TF) is determined as the ratio of heavy metal concentration in plant shoot to that in root.

1.4. Plant growth and root activity analysis

Fresh weight (fw) and dry weight of plant were measured by electronic balance as an indicator of plant growth. The water content (C_w , %) was determined on the basis of Eq. (1):

$$C_w = \frac{W_f - W_d}{W_f} \times 100\% \quad (1)$$

where, W_f (g) and W_d (g) refer to fresh weight and dry weight of plant, respectively.

Root activity (RA) was determined by the triphenyl tetrazolium chloride (TTC) method (Clemensson-Lindell, 1994). Fresh roots (about 0.5 g) were immersed in beakers with the addition of 10 mL of 0.4% TTC and phosphate buffer (1/15 mol/L, pH 7.0). After incubation at 37°C for 2 hr in the dark, 2 mL of 1 mol/L H_2SO_4 was added to terminate the reaction. Then the samples

were ground homogeneously in 4 mL of 95% (V/V) ethyl acetate to extract the reduced triphenyl tetrazolium formazan (TTF). The extracts were transferred into a graduated test tube and the residues were washed three times with the extract buffer. The absorbance of the color was recorded at 485 nm and RA was calculated as $\text{mg TTF} / (\text{g fw} \cdot \text{hr})$.

1.5. Physiological assay

Extraction of chlorophyll and carotenoid was carried out according to the method of Lichtenthaler (1987). About 0.5 g frozen leaves were ground homogeneously with 80% chilled acetone in the dark and centrifuged at 2000 r/min for 10 min. Then chlorophyll and carotenoid content were read by ultraviolet spectrophotometer (UV-2550, Shimadzu, Kyoto, Japan) on the supernatant at 470, 646 and 663 nm. Chlorophyll and carotenoid content were expressed as mg/g fw.

Following the method of Chaoui et al. (1997), the malondialdehyde (MDA) content was determined by the 2-thiobarbituric acid (TBA) reactive metabolites. Leaf and root samples (about 1 g fw, respectively) were ground homogeneously in 10 mL of 10% (W/V) trichloroacetic acid (TCA). The extract was centrifuged at 10,000 r/min, 4°C for 10 min. Absorbing 2 mL of the supernatant, 3 mL of 0.5% TBA (made in 10% TCA) was added. After incubation at 95°C for 30 min, the mixture was ice-cooled quickly and then centrifuged at 10,000 r/min, 4°C for 15 min. The absorbance of the supernatant was measured at 532 nm and corrected for non-specific turbidity at 600 nm. The MDA content was calculated as nmol/g fw.

Soluble protein was assayed by the binding of Coomassie Brilliant Blue G-250 method (Bradford, 1976). Leaf and root samples (about 0.5 g fw, respectively) were treated with 3 mL of solution containing 50 mmol/L Tris-HCl (pH 7.8), 0.5 mmol/L MgCl_2 , 1 mmol/L EDTA and 1 mmol/L dithiothreitol in ice precooled mortar. The mixture was homogenized under ice-bath condition and centrifuged at 10,000 r/min, 4°C for 20 min. Then the supernatant (0.1 mL) was mixed with 0.9 mL of Tris buffer solution and 5 mL of Coomassie Brilliant Blue G-250 staining solution (Philadelphia, USA) completely. After being kept for 2 min, the absorbance of the extract was measured at 595 nm corresponding with the bovine serum albumin calibration curve. The soluble protein content was measured as mg/g fw.

Frozen leaf samples (0.5 g fw) were homogenized under chilled condition in the buffer: 50 mmol/L, pH 7.8 phosphate buffer containing 0.1% (V/V) Triton X-100 and 1% (W/V) polyvinylpyrrolidone for extracting SOD and POD; 0.9% (W/V) NaCl was used as an extraction buffer for CAT extraction. Then the homogenate was centrifuged at 15,000 r/min, 4°C for 20 min and the supernatant was used for determining enzyme activities, which were processed at 25°C. SOD activity was assayed by photochemical nitroblue tetrazolium (NBT) method. Photoreduction of NBT (formation of purple formazan) at 50% inhibition was recorded at 560 nm using 3 mL of mixture containing 50 mmol/L phosphate buffer (pH 7.8), 750 $\mu\text{mol/L}$ NBT, 130 mmol/L methionine, 100 $\mu\text{mol/L}$ EDTA- Na_2 and 20 $\mu\text{mol/L}$ riboflavin. POD activity was determined using Guaiacol method by monitoring the increase in absorption at 470 nm as a consequence of guaiacol oxidation in a reaction solution (3 mL final volume) consisting of 50 mmol/L

phosphate buffer (pH 7.0), 0.3% H_2O_2 , 0.2% guaiacol and 0.5 mL of enzyme extract. For CAT activity determination, the H_2O_2 reduction method was used to measure the decrease in absorbance at 240 nm due to H_2O_2 consumption (Aebi, 1984). The reaction mixture contained 50 mmol/L phosphate buffer (pH 7.0), 0.1 mol/L H_2O_2 and 0.5 mL of leaf enzyme extract solution in a 3 mL volume. These enzyme activities were expressed as U/mg protein, $\text{OD}_{470} / (\text{g fw} \cdot \text{min})$ and $\text{OD}_{240} / (\text{g fw} \cdot \text{min})$, respectively, where 1 U meant enzyme content when the relative inhibition ratio of NBT photoreduction reached 50%.

1.6. Data analysis

The results from a representative experiment were presented as mean values \pm S.E. of three replications. Graphical work was carried out using Origin v.8.0 (USA). Statistical significance was conducted by t-test at a probability level of $p < 0.05$.

2. Results

2.1. Cadmium accumulation and translocation

In the hydroponic condition, Cd accumulation in ramie roots and shoots exhibited a linear increase in response to elevated Cd in the medium as shown in Table 1. Cd accumulated in plant root was approximately 9–10 times that in the aerial part. It appeared to change significantly from Cd₂ to Cd₅ group ($p < 0.05$), with Cd uptake from 268.0 to 370.8 mg/kg dw in root and 25.2 to 35.6 mg/kg dw in shoot. The presence of citric and oxalic acids promoted Cd uptake in plant aerial and underground parts. Especially, the low dose of citric acid dramatically increased Cd uptake by 26.7% in root and 1-fold in shoot ($p < 0.01$). A slight increase in Cd transport from roots to shoots was observed, ranging from 9.4% to 11.0%. Compared to Cd₁₀ treatment, the low dose of citric and oxalic acids significantly increased TF from 11.0% in the control to 17.8% and 18.3% ($p < 0.05$), respectively.

2.2. Plant growth attributes and RA

Plants showed slight symptoms of Cd phytotoxicity, as seen from trivial chlorosis in young leaves and browned symptom in root tips after exposure to Cd for 1 week. Plant growth retardation was observed at all Cd levels with a reduction in both fresh biomass and dry biomass, as well as shoot/root ratio (Table 2). Particularly fresh biomass and dry biomass at 2 mg/L of Cd level were declined by 30.5% and 46.7%, respectively. Compared to Cd₁₀ treatment, the citric and oxalic acid addition alleviated the inhibition to plant growth, as seen from the increase in fresh biomass and dry biomass of plant tissues. The low level of citric acid enhanced plant fresh biomass and dry biomass by 44.6% and 74.4%, respectively, which were more distinct than the high one and two levels of oxalic acid. The water content increased from 74.4% in the control to 84.4% at 5 mg/L of Cd level, and decreased to 84.0% at highest Cd level. Citric acid decreased it to 80.8%, whereas oxalic acid slightly elevated it to 86.2%. The changes are not significant.

RA is a reliable indicator of root respiration and mineral substance absorption capability. As shown in Fig. 1, RA reached

Table 1 – Effects of Cd, citric and oxalic acids on Cd uptake and translocation in ramie (mean \pm SE, $n = 3$).

Treatment	Cd concentration (mg/kg dw)		Root/shoot ratio of Cd concentration	TF value (%)
	Shoot	Root		
Cd ₀	ND	ND	–	–
Cd ₂	25.2 \pm 2.6	268.0 \pm 24.3	10.6	9.4
Cd ₅	35.6 \pm 4.2*	370.8 \pm 36.5*	10.4	9.6
Cd ₁₀	41.2 \pm 4.0	374.4 \pm 35.3	9.1	11.0
Cd ₁₀ + CA ₁	84.2 \pm 8.7**	474.2 \pm 45.8**	5.6	17.8*
Cd ₁₀ + CA ₂	56.6 \pm 5.8	409.0 \pm 37.5*	7.2	13.8
Cd ₁₀ + OA ₁	84.0 \pm 8.0**	460.2 \pm 43.8**	5.5	18.3*
Cd ₁₀ + OA ₂	53.0 \pm 4.7	389.4 \pm 34.7	7.4	13.6

Cd₀: without addition of Cd in nutrient solution; Cd₂, Cd₅, Cd₁₀: 2, 5, 10 mg/L Cd concentration; Cd₁₀ + CA₁: 10 mg/L Cd concentration + 1.5 mmol/L citric acid; Cd₁₀ + CA₂: 10 mg/L Cd concentration + 4 mmol/L citric acid; Cd₁₀ + OA₁: 10 mg/L Cd concentration + 3 mmol/L oxalic acid; Cd₁₀ + OA₂: 10 mg/L Cd concentration + 9 mmol/L oxalic acid.

ND: not detectable.

TF: translocation factor, the ratio of heavy metal concentration in plant shoot to that in root.

* and ** indicate significant differences in parameters between the control group and the experimental group at $p < 0.05$ and $p < 0.01$, respectively.

the highest value in the control (4.1 mg TTF / (g fw · hr)) and decreased progressively to 62.2%, 42.2% and 12.5% of the control with Cd level increasing. It was improved significantly to 15.2% of the control in the presence of low citric acid level ($p < 0.05$). But no visible changes occurred in oxalic acid treatment.

2.3. Chlorophyll and carotenoid content

Chlorophyll is regarded as a key indicator of plant photosynthetic capacity. Carotenoid, a cell endogenous antioxidant, plays an important role in scavenging ROS and preventing membrane lipid peroxidation. Chlorophyll content reached maximum (7.8 mg/g fw, 2.3 times of the control) at 5 mg/L of Cd level and declined sharply (5.3 mg/g fw) at highest Cd level. However, carotenoid content increased with Cd levels, reaching maximum (1.3 mg/g fw, 4.5 times of the control) at 10 mg/L of Cd level (Fig. 2). The low dose of citric acid (1.5 mmol/L) significantly increased chlorophyll and carotenoid content by 17.9% and 35.7% ($p < 0.05$), respectively, which exhibited a better mitigating effect on photosynthetic pigments than the high dose and two doses of oxalic acid.

2.4. Lipid peroxidation

The extent of lipid peroxidation in plant leaves and roots was estimated by determining MDA content. The much higher MDA content of ramie leaves revealed severer lipid peroxidation emerging in leaves. There was a dose-dependent increase

in MDA content with a maximum of 57.1% increase in leaf and 72.1% in root at 10 mg/L of Cd level compared to the control (Fig. 3). Citric and oxalic acids ameliorated lipid peroxidation with MDA content decrease comparably in contrast to Cd₁₀ treatment alone, especially the high citric acid level with 26.2% and 29.4% reduction in leaf and root, respectively. Citric acid exhibited a better alleviation on lipid peroxidation than oxalic acid, and the high dose of acid was better than the low one.

2.5. Soluble protein

Soluble protein content increased significantly by 31.8% in leaf and 76.3% in root at 5 mg/L of Cd level compared to the control ($p < 0.05$). Then it declined slightly by 7.3% in leaf and sharply by 46.2% (the minimum value 30.8 mg/g fw) in root at highest Cd level probably due to protein degradation (Fig. 4). The presence of citric acid resulted in enhancement of soluble protein content with an average of 2.9, 6.4 mg/g increment in leaf and root, respectively. While the high dose of oxalic acid declined the content by 3.1 mg/g in leaf.

2.6. Antioxidative enzymes

The SOD activity increased dramatically at low Cd level in response to Cd stress and this rise trend became gentle under higher Cd concentrations (Fig. 5). It reached 1.8-fold of non-Cd treatment at highest Cd level. The presence of citric and oxalic acids further enhanced the SOD activity. Especially the low citric

Table 2 – Effects of Cd, citric and oxalic acids on fresh biomass, dry biomass and water content of ramie (mean \pm SE, $n = 3$).

Treatment	Fresh biomass (g/plant)			Dry biomass (g/plant)			Water content (%)
	Shoot	Root	Shoot/root ratio	Shoot	Root	Shoot/root ratio	
Cd ₀	17.62 \pm 1.61	9.25 \pm 0.95	1.90	4.83 \pm 0.52	2.06 \pm 0.23	2.34	74.4
Cd ₂	11.06 \pm 1.27	7.61 \pm 0.69	1.45	2.41 \pm 0.21	1.26 \pm 1.39	1.91	80.3
Cd ₅	8.28 \pm 0.76	7.93 \pm 0.72	1.04	1.42 \pm 0.15	1.11 \pm 0.09	1.28	84.4
Cd ₁₀	7.72 \pm 0.73	5.50 \pm 0.58	1.40	1.29 \pm 0.14	0.82 \pm 0.07	1.57	84.0
Cd ₁₀ + CA ₁	10.72 \pm 1.06	8.40 \pm 0.83	1.28	2.36 \pm 0.25	1.32 \pm 0.14	1.79	80.8
Cd ₁₀ + CA ₂	9.10 \pm 0.98	8.38 \pm 0.81	1.09	1.76 \pm 0.16	1.27 \pm 0.12	1.39	82.7
Cd ₁₀ + OA ₁	9.86 \pm 0.92	6.48 \pm 0.65	1.52	1.56 \pm 0.13	0.95 \pm 0.08	1.64	84.6
Cd ₁₀ + OA ₂	8.72 \pm 0.85	6.78 \pm 0.66	1.29	1.30 \pm 0.12	0.84 \pm 0.06	1.55	86.2




Fig. 1 – Root activity of ramie exposed to different levels of Cd, citric and oxalic acid application at the highest Cd level. Vertical bars means the standard error of mean values ($n = 3$). * and ** indicate significant differences in parameters between the control group and the experimental group at $p < 0.05$ and $p < 0.01$, respectively.

acid level improved SOD activity by 7.8%, more significantly than the high level as well as two levels of oxalic acid. Activities of POD and CAT decreased gradually with elevated Cd concentration and reached a minimum at highest Cd level (81.6% and 64.5% of its control, respectively). Changes in CAT activity followed the similar pattern as POD activity. A considerable enhancement of the two enzyme activities was observed in the presence of organic acids, which facilitated the efficiency of scavenging ROS and alleviated oxidative stress from Cd. Particularly, low citric acid level exhibited a better alleviation (7.5% increase in POD activity and 7.4% in CAT activity) than the high level as well as two levels of oxalic acid.

3. Discussion

3.1. Prompting effects on Cd uptake and translocation by citric and oxalic acids

It is meaningful to study the accumulation and transport mechanism of metals in plant for the high efficiency of phytoremediation. Cd content in plant tissues increased in a

dose-dependent manner, which are in accordance with the results reported by Fidalgo et al. (2011). Compared to the aerial tissues, ramie showed a tendency to accumulate a higher amount of Cd in roots that were primary sites for metal access, as a natural protective response to lessen Cd toxicity in plant aboveground tissues especially leaves. Therefore, the translocation factor of Cd from roots to shoots remained at a relatively low value. Citric acid at 1.5 mmol/L and oxalic acid at 3 mmol/L significantly increased Cd uptake to both 2-fold in the shoots and 1.3 and 1.2-folds in the roots of the controls, respectively. They also improved the TF by 61.8% and 66.4%, respectively. Citric and oxalic acids promoted Cd uptake and transport, in consistent with the findings of Duarte et al. (2007), who reported enhanced Cd uptake and translocation by *Halimione portulacoides* in the presence of citric acid.

The above fact was probably due to the following reasons. Firstly, Cd solubility and availability were increased by the increased pH and the release of strong organic ligands from organic acids into the plant–solution interface, which facilitated Cd uptake by ramie roots and translocation to the shoots. Secondly, they activated ATPases in the plasma membrane resulting in alteration of ions transport in charge of metal




Fig. 2 – Total chlorophyll and carotenoid content in leaves of ramie exposed to different levels of Cd, citric and oxalic acid application at the highest Cd level. Vertical bars refer to the standard error of mean values ($n = 3$). * and ** indicate significant differences in parameters between the control group and the experimental group at $p < 0.05$ and $p < 0.01$, respectively.




Fig. 3 – Malondialdehyde (MDA) content in leaves and roots of ramie exposed to different levels of Cd, citric and oxalic acid application at the highest Cd level. Vertical bars refer to the standard error of mean values ($n = 3$). * and ** indicate significant differences in parameters between the control group and the experimental group at $p < 0.05$ and $p < 0.01$, respectively.

translocation. Furthermore, organic acid chelating with heavy metal should be viewed as an essential process of efficiently participating in metal long-distance translocation in the xylem and avoiding the toxicity of free metal ions in plants (Evangelou et al., 2006; Montargès-Pelletier et al., 2008; Wei et al., 2009; Ghnaya et al., 2013). The chelant-metal complex might enter into plant tissues through breaking in the root endodermis and Casparian strips, and subsequently translocating upward to the shoots with the assistance of transpiration.




Fig. 4 – Soluble protein content in leaves and roots of ramie exposed to different levels of Cd, citric and oxalic acid application at the highest Cd level. Vertical bars refer to the standard error of mean values ($n = 3$). * and ** indicate significant differences in parameters between the control group and the experimental group at $p < 0.05$ and $p < 0.01$, respectively.




Fig. 5 – Superoxide dismutase (SOD), Peroxidase (POD) and Catalase (CAT) activities in leaves of ramie exposed to different levels of Cd, citric and oxalic acid application at the highest Cd level. Vertical bars means the standard error of mean values ($n = 3$).

3.2. Effects of citric and oxalic acids on plant physiological properties

The present study was carried out in hydroponics for the reason that the medium provided a homogeneous condition in favor of quick identification of Cd-induced plant physiological characters. Results showed that ramie was able to survive moderate Cd oxidative injury with some repairment of self-defense system and 10 mg/L was regarded as its tolerance threshold to Cd. Citric acid at 1.5 mmol/L could play a remarkable role in regulating Cd-induced physiological alterations in ramie.

A better comprehension of heavy metal toxicity on physiological metabolism contributes to investigating the approaches for Cd tolerance enhancement. An increasing metal sequestration, tissue tolerance and antioxidative

response determine the ability of plant to defy metal destructive damage (Jin et al., 2008). Heavy metal might interfere with numerous biochemical and physiological processes including respiration, nutrient uptake, photosynthetic system, protein metabolism and activities of antioxidative enzymes (Shamsi et al., 2008; Daud et al., 2009; Rai et al., 2013). Reduced plant biomass could reflect an irreversible plant growth inhibition (Srivastava et al., 2004), which appeared to be correlated with the decrease in RA under water stress induced by Cd stress. Reduced RA in turn resulted in weaker root respiration, less ATP production and nutrient elements absorption, as well as the reduction in POD activity involving in plant growth and development processes. The gradually brownish tissue and the sharp decrease of RA exhibited its sensibility to Cd toxicity. Cd stress resulted in a distinct decrease of chlorophyll and soluble protein synthesis. Particularly, the higher Cd concentration not only caused a dramatic breakdown of chlorophyll by inhibiting protochlorophyllide reductase and the photosynthetic electron transport (Aibibu et al., 2010), but also resulted in protein metabolism degradation in excess of Cd toxicity beyond plant's tolerance extreme. The changes of chlorophyll content and soluble protein content in this study coincided with the findings of Liu et al. (2007).

Antioxidative enzymes and certain metabolites play a crucial role in adaptation and ultimate survival of plant to Cd oxidative damage. With the Cd stress rising, there was an increase in the activity of SOD and a decline in the activities of POD and CAT, in line with the results of Najeeb et al. (2011). Enhanced relative water content (generally as the osmotic influence of abiotic stress), photosynthetic pigments, protein synthesis and SOD activity could be viewed as an adaptive regulation in response to Cd stress. Especially the SOD activity in the antioxidant system increased progressively with increasing Cd level, and acted as the first defense line of scavenging ROS to avoid excess oxidative impairment in response to the stimulation of lipid peroxidation. However, POD and CAT activities showed similar descending tendency probably due to the inhibition of enzyme synthesis or an alteration in the assemblage of enzyme subunits. Furthermore, inhibition of POD and CAT synthesis limited the efficiency of eliminating H_2O_2 produced in chloroplasts, which would lead to membrane lipid peroxidation demonstrated by the continuous elevated MDA content (an indicator of oxidative injury), implying that oxidative stress from Cd (probably the deleterious effect of H_2O_2) did exist in plant. Generally, the lipid peroxidation exhibited a concentration-dependent increasing trend. Particularly, it was relatively severer in ramie leaves than roots (Fig. 3). This could be explained by the following scenario: the generation of $NADP^+$ (a key electron acceptor in photosynthesis) decreased due to damaged CO_2 assimilation resulting from stomatal closure under Cd stress in leaves, further the lacking of $NADP^+$ with continuous light reaction form O_2^- and subsequent H_2O_2 by facilitating the electron transfer from PS-I to O_2 (Demiral and Türkan, 2005; Liu et al., 2007).

In the presence of organic acids, all antioxidative enzyme activities were enhanced and lipid peroxidation was mitigated compared with the controls, which suggested that organic acids might have interfered in combating Cd-induced oxidative stress in plant as a consequence of the less toxicity of organic acid-Cd chelates. Citric acid posed a more effective

influence due to its stronger binding affinity for Cd than oxalic acid. The contents of chlorophyll, carotenoid and soluble protein were also elevated owing to the alleviation of Cd oxidative damage. This could be explained by the prevention of free metal ion transmission in the cytosol. Cd damaged plant cells through cytoplasmic shrinkage and metal deposition and citric acid restored structure and shape of cells and eliminated plasmolysis (Najeeb et al., 2011). Thus the insight was proposed that ramie was able to survive more easily under high Cd stress, and exerted stronger defense in the presence of organic acids especially the citric acid, without showing visible phytotoxicity symptoms, which had prospect to be a useful strategy for the phytoremediation of Cd contaminated area, particularly the Cd polluted farmland concerning safety grain production.

Cd has previously been reported to induce phytochelatin (PC) synthesis using GSH as its precursor in several plant species (Srivastava et al., 2004; Vurro et al., 2011) particularly in root as a detoxify mechanism. In the subsequent work, we would like to investigate the interactions between PC induction and plant detoxification under abiotic Cd stress and the role of organic acids in this process to examine plant physiological mechanism more deeply (Mishra et al., 2006; Seth et al., 2008).

4. Conclusions

The results obtained from the present study demonstrated that exogenous citric and oxalic acids had the capability of facilitating Cd uptake and transport, and alleviating the physiological toxicity from free metal ions. As a Cd tolerance plant, ramie accumulated more Cd in root tissue than that in shoot tissue in order to help plant survive Cd toxicity. High Cd level reduced plant biomass, root activity, chlorophyll content, synthesis of protein, antioxidative enzyme activities (POD and CAT) and enlarged lipid peroxidation. Furthermore, citric and oxalic acids acted as a ligand to chelate with Cd and entered in the form of organic acid-Cd chelation, which efficiently alleviated the physiological toxicity from free metal ions. Taken together, as a non-edible crop and a pioneering plant species of abandoned mining area, ramie in combination with citric acid has promise to be applied in the phytoremediation of moderate Cd contaminated site due to the increased plant biomass, Cd uptake and endurance.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No. 41271332), the Natural Science Foundation of Hunan province (No. 11JJ2031) and the Science and Technology Planning Project of Hunan Province, China (No. 2012SK2021). The authors are grateful to the Key Laboratory of Environmental Biology and Pollution Control (Hunan University) for creating a good research place with an advanced technical apparatus. We also thank Mrs. X. Wang from College of Resources and Environmental Science (Hunan Normal University) for the offer of research fellowship and language assistance.

REFERENCES

- Aeby, H., 1984. Catalases. In: Bergmeyer, H.U. (Ed.), *Methods of Enzymatic Analysis* vol. 2. Academic Press, New York, pp. 673–684.
- Aibibu, N.A., Liu, Y.G., Zeng, G.M., Wang, X., Chen, B.B., Song, H.X., et al., 2010. Cadmium accumulation in *Vetiveria zizanioides* and its effects on growth, physiological and biochemical characters. *Bioresour. Technol.* 101 (16), 6297–6303.
- Bhargava, A., Carmona, F.F., Bhargava, M., Srivastava, S., 2012. Approaches for enhanced phytoextraction of heavy metals. *J. Environ. Manag.* 105, 103–120.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Anal. Biochem.* 72 (1–2), 248–254.
- Chaoui, A., Mazhoudi, S., Ghorbal, M.H., Ferjani, E.E., 1997. Cadmium and zinc induction of lipid peroxidation and effects on antioxidant enzyme activities in bean (*Phaseolus vulgaris* L.). *Plant Sci.* 127 (2), 139–147.
- Clabeaux, B.L., Navarro, D.A.G., Aga, D.S., Bisson, M.A., 2011. Cd tolerance and accumulation in the aquatic macrophyte, *Chara australis*: potential use for Charophytes in phytoremediation. *Environ. Sci. Technol.* 45 (12), 5332–5338.
- Clemensson-Lindell, A., 1994. Triphenyltetrazolium chloride as an indicator of fine-root vitality and environmental stress in coniferous forest stands: applications and limitations. *Plant Soil* 159 (2), 297–300.
- Daud, M.K., Sun, Y.Q., Dawood, M., Hayat, Y., Variath, M.T., Wu, Y.X., et al., 2009. Cadmium-induced functional and ultrastructural alterations in roots of two transgenic cotton cultivars. *J. Hazard. Mater.* 161 (1), 463–473.
- Demiral, T., Türkan, I., 2005. Comparative lipid peroxidation, antioxidant defense systems and proline content in roots of two rice cultivars differing in salt tolerance. *Environ. Exp. Bot.* 53 (3), 247–257.
- Duarte, B., Delgado, M., Cacador, I., 2007. The role of citric acid in cadmium and nickel uptake and translocation, in *Halimione portulacoides*. *Chemosphere* 69 (5), 836–840.
- Edreva, A., 2005. Generation and scavenging of reactive oxygen species in chloroplasts: a submolecular approach. *Agric. Ecosyst. Environ.* 106 (2–3), 119–133.
- Evangelou, M.W.H., Ebel, M., Schaeffer, A., 2006. Evaluation of the effect of small organic acids on phytoextraction of Cu and Pb from soil with tobacco *Nicotiana tabacum*. *Chemosphere* 63 (6), 996–1004.
- Fernández, R., Bertrand, A., Reis, R., Mourato, M.P., Martins, L.L., González, A., 2013. Growth and physiological responses to cadmium stress of two populations of *Dittrichia viscosa* (L.) Greuter. *J. Hazard. Mater.* 244–245, 555–562.
- Fidalgo, F., Freitas, R., Ferreira, R., Pessoa, A.M., Teixeira, J., 2011. *Solanum nigrum* L. antioxidant defence system isozymes are regulated transcriptionally and posttranslationally in Cd-induced stress. *Environ. Exp. Bot.* 72 (2), 312–319.
- Ghnaya, T., Zaier, H., Baioui, R., Sghaier, S., Lucchini, G., Sacchi, G. A., et al., 2013. Implication of organic acids in the long-distance transport and the accumulation of lead in *Sesuvium portulacastrum* and *Brassica juncea*. *Chemosphere* 90 (4), 1449–1454.
- Irtelli, B., Navari-Izzo, F., 2006. Influence of sodium nitrilotriacetate (NTA) and citric acid on phenolic and organic acids in *Brassica juncea* grown in excess of cadmium. *Chemosphere* 65 (8), 1348–1354.
- Jean, L., Bordas, F., Gautier-Moussard, C., Vernay, P., Hitmi, A., Bollinger, J.C., 2008. Effect of citric acid and EDTA on chromium and nickel uptake and translocation by *Datura innoxia*. *Environ. Pollut.* 153 (3), 555–563.
- Jin, X.F., Yang, X.E., Islam, E., Liu, D., Mahmood, Q., 2008. Effects of cadmium on ultrastructure and antioxidative defense system in hyperaccumulator and non-hyperaccumulator ecotypes of *Sedum alfredii* Hance. *J. Hazard. Mater.* 156 (1–3), 387–397.
- Lei, M., Zeng, M., Wang, L.H., Williams, P.N., Sun, G.X., 2010. Arsenic, lead and cadmium pollution in rice from Hunan markets and contaminated areas and their health risk assessment. *Acta Sci. Circumst.* 30 (11), 2314–2320.
- Li, X., Zhao, M.X., Guo, L.P., Huang, L.Q., 2012. Effect of cadmium on photosynthetic pigments, lipid peroxidation, antioxidant, and artemisinin in hydroponically grown *Artemisia annua*. *J. Environ. Sci.* 24 (8), 1511–1518.
- Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. In: Baltscheffsky, M. (Ed.), *Current Research in Photosynthesis*. Kluwer Academic Publishers, Dordrecht, pp. 471–474.
- Liu, Y.G., Wang, X., Zeng, G.M., Qu, D., Gu, J.J., Zhou, M., et al., 2007. Cadmium-induced oxidative stress and response of the ascorbate–glutathione cycle in *Beckermia nivea* (L.) Gaud. *Chemosphere* 69 (1), 99–107.
- Liu, H.J., Zhang, J.L., Christie, P., Zhang, F., 2008. Influence of iron plaque on uptake and accumulation of Cd by rice (*Oryza sativa* L.) seedlings grown in soil. *Sci. Total Environ.* 394 (2–3), 361–368.
- López-Millán, A.F., Sagardoy, R., Solanas, M., Abadía, A., Abadía, J., 2009. Cadmium toxicity in tomato (*Lycopersicon esculentum*) plants grown in hydroponics. *Environ. Exp. Bot.* 65 (2–3), 376–385.
- Meighan, M.M., Fenus, T., Karey, E., MacNeil, J., 2011. The impact of EDTA on the rate of accumulation and root/shoot partitioning of cadmium in mature dwarf sunflowers. *Chemosphere* 83 (11), 1539–1545.
- Mishra, S., Srivastava, S., Tripathi, R.D., Govindarajan, R., Kuriakose, S.V., Prasad, M.N.V., 2006. Phytochelatin synthesis and response of antioxidant during cadmium stress in *Bacopa monnieri* L. *Plant Physiol. Biochem.* 44 (1), 25–37.
- Montargès-Pelletier, E., Chardot, V., Echevarria, G., Michot, L.J., Bauer, A., Morel, J.L., 2008. Identification of nickel chelators in three hyperaccumulating plants: an X-ray spectroscopic study. *Phytochemistry* 69 (8), 1695–1709.
- Najeeb, U., Xu, L., Ali, S., Jilani, G., Gong, H.J., Shen, W., et al., 2009. Citric acid enhances the phytoextraction of manganese and plant growth by alleviating the ultrastructural damages in *Juncus effusus* L. *J. Hazard. Mater.* 170 (2–3), 1156–1163.
- Najeeb, U., Jilani, G., Ali, S., Sarwar, M., Xu, L., Zhou, W.J., 2011. Insights into cadmium induced physiological and ultra-structural disorders in *Juncus effusus* L. and its remediation through exogenous citric acid. *J. Hazard. Mater.* 186 (1), 565–574.
- Pérez-Esteban, J., Escolástico, C., Moliner, A., Masaguer, A., 2013. Chemical speciation and mobilization of copper and zinc in naturally contaminated mine soils with citric and tartaric acids. *Chemosphere* 90 (2), 276–283.
- Rai, U.N., Singh, N.K., Upadhyay, A.K., Verma, S., 2013. Chromate tolerance and accumulation in *Chlorella vulgaris* L.: role of antioxidant enzymes and biochemical changes in detoxification of metals. *Bioresour. Technol.* 136, 604–609.
- Seth, C.S., Chaturvedi, P.K., Misra, V., 2008. The role of phytochelators and antioxidants in tolerance to Cd accumulation in *Brassica juncea* L. *Ecotoxicol. Environ. Saf.* 71 (1), 76–85.
- Shamsi, I.H., Jilani, G., Zhang, G.P., Wei, K., 2008. Cadmium stress tolerance through potassium nutrition in soybean. *Asian J. Chem.* 20 (2), 1099–1108.
- Srivastava, S., Tripathi, R.D., Dwivedi, U.N., 2004. Synthesis of phytochelators and modulation of antioxidants in response to cadmium stress in *Cuscuta reflexa* — an angiospermic parasite. *J. Plant Physiol.* 161 (6), 665–674.

- Vurro, E., Ruotolo, R., Ottonello, S., Elviri, L., Maffini, M., Falasca, G., et al., 2011. Phytochelatins govern zinc/copper homeostasis and cadmium detoxification in *Cuscuta campestris* parasitizing *Daucus carota*. *Environ. Exp. Bot.* 72 (1), 26–33.
- Wang, X., Liu, Y.G., Zeng, G.M., Chai, L.Y., Song, X.C., Min, Z.Y., et al., 2008. Subcellular distribution and chemical forms of cadmium in *Beckmannia nivea* (L.) Gaud. *Environ. Exp. Bot.* 62 (3), 389–395.
- Wang, X., Ma, L.Q., Rathinasabapathi, B., Cai, Y., Liu, Y.G., Zeng, G.M., 2011. Mechanisms of efficient arsenite uptake by arsenic hyperaccumulator *Pteris vittata*. *Environ. Sci. Technol.* 45 (22), 9719–9725.
- Wei, W., Wang, Y., Wei, Z.G., Zhao, H.Y., Li, H.X., Hu, F., 2009. Roles of organic acids and nitrate in the long-distance transport of cobalt in xylem saps of *Alyssum murale* and *Trifolium subterraneum*. *Biol. Trace Elem. Res.* 131 (2), 165–176.
- Xia, W.B., Gao, H., Wang, X.H., Zhou, C.H., Liu, Y.G., Fan, T., et al., 2009. Application of EDTA decontamination on soils affected by mining activities and impact of treatment on the geochemical partition of metal contaminants. *J. Hazard. Mater.* 164 (2–3), 936–940.
- Yang, L., Luo, C.L., Liu, Y., Quan, L.T., Chen, Y.H., Shen, Z.G., 2013. Residual effects of EDDS leachates on plants during EDDS-assisted phytoremediation of copper contaminated soil. *Sci. Total Environ.* 444, 263–270.



Editorial Board of Journal of Environmental Sciences

Editor-in-Chief

Hongxiao Tang Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, China

Associate Editors-in-Chief

Jiuhui Qu Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, China
Shu Tao Peking University, China
Nigel Bell Imperial College London, United Kingdom
Po-Keung Wong The Chinese University of Hong Kong, Hong Kong, China

Editorial Board

Aquatic environment

Baoyu Gao
Shandong University, China
Maohong Fan
University of Wyoming, USA
Chihpin Huang
National Chiao Tung University
Taiwan, China
Ng Wun Jern
Nanyang Environment &
Water Research Institute, Singapore
Clark C. K. Liu
University of Hawaii at Manoa, USA
Hokyoung Shon
University of Technology, Sydney, Australia
Zijian Wang
Research Center for Eco-Environmental Sciences,
Chinese Academy of Sciences, China
Zhiwu Wang
The Ohio State University, USA
Yuxiang Wang
Queen's University, Canada
Min Yang
Research Center for Eco-Environmental Sciences,
Chinese Academy of Sciences, China
Zhifeng Yang
Beijing Normal University, China
Han-Qing Yu
University of Science & Technology of China

Terrestrial environment

Christopher Anderson
Massey University, New Zealand
Zucong Cai
Nanjing Normal University, China
Xinbin Feng
Institute of Geochemistry,
Chinese Academy of Sciences, China
Hongqing Hu
Huazhong Agricultural University, China
Kin-Che Lam
The Chinese University of Hong Kong
Hong Kong, China
Erwin Klumpp
Research Centre Juelich, Agrosphere Institute
Germany
Peijun Li
Institute of Applied Ecology,
Chinese Academy of Sciences, China

Michael Schlöter

German Research Center for Environmental Health
Germany
Xuejun Wang
Peking University, China
Lizhong Zhu
Zhejiang University, China

Atmospheric environment

Jianmin Chen
Fudan University, China
Abdelwahid Mellouki
Centre National de la Recherche Scientifique
France
Yujing Mu
Research Center for Eco-Environmental Sciences,
Chinese Academy of Sciences, China
Min Shao
Peking University, China
James Jay Schauer
University of Wisconsin-Madison, USA
Yuesi Wang
Institute of Atmospheric Physics,
Chinese Academy of Sciences, China
Xin Yang
University of Cambridge, UK

Environmental biology

Yong Cai
Florida International University, USA
Henner Hollert
RWTH Aachen University, Germany
Jae-Seong Lee
Sungkyunkwan University, South Korea
Christopher Rensing
University of Copenhagen, Denmark
Bojan Sedmak
National Institute of Biology, Slovenia
Lirong Song
Institute of Hydrobiology,
Chinese Academy of Sciences, China
Chunxia Wang
National Natural Science Foundation of China
Gehong Wei
Northwest A & F University, China
Daqiang Yin
Tongji University, China
Zhongtang Yu
The Ohio State University, USA

Environmental toxicology and health

Jingwen Chen
Dalian University of Technology, China
Jianying Hu
Peking University, China
Guibin Jiang
Research Center for Eco-Environmental Sciences,
Chinese Academy of Sciences, China
Sijin Liu
Research Center for Eco-Environmental Sciences,
Chinese Academy of Sciences, China
Tsuyoshi Nakanishi
Gifu Pharmaceutical University, Japan
Willie Peijnenburg
University of Leiden, The Netherlands
Bingsheng Zhou
Institute of Hydrobiology,
Chinese Academy of Sciences, China

Environmental catalysis and materials

Hong He
Research Center for Eco-Environmental Sciences,
Chinese Academy of Sciences, China
Junhua Li
Tsinghua University, China
Wenfeng Shangguan
Shanghai Jiao Tong University, China
Yasutake Teraoka
Kyushu University, Japan
Ralph T. Yang
University of Michigan, USA

Environmental analysis and method

Zongwei Cai
Hong Kong Baptist University,
Hong Kong, China
Jiping Chen
Dalian Institute of Chemical Physics,
Chinese Academy of Sciences, China
Minghui Zheng
Research Center for Eco-Environmental Sciences,
Chinese Academy of Sciences, China

Municipal solid waste and green chemistry

Pinjing He
Tongji University, China
Environmental ecology
Rusong Wang
Research Center for Eco-Environmental Sciences,
Chinese Academy of Sciences, China

Editorial office staff

Managing editor Qingcai Feng
Editors Zixuan Wang Suqin Liu Zhengang Mao
English editor Catherine Rice (USA)

JOURNAL OF ENVIRONMENTAL SCIENCES

环境科学学报(英文版)
(<http://www.jesc.ac.cn>)

Aims and scope

Journal of Environmental Sciences is an international academic journal supervised by Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences. The journal publishes original, peer-reviewed innovative research and valuable findings in environmental sciences. The types of articles published are research article, critical review, rapid communications, and special issues.

The scope of the journal embraces the treatment processes for natural groundwater, municipal, agricultural and industrial water and wastewaters; physical and chemical methods for limitation of pollutants emission into the atmospheric environment; chemical and biological and phytoremediation of contaminated soil; fate and transport of pollutants in environments; toxicological effects of terrorist chemical release on the natural environment and human health; development of environmental catalysts and materials.

For subscription to electronic edition

Elsevier is responsible for subscription of the journal. Please subscribe to the journal via <http://www.elsevier.com/locate/jes>.

For subscription to print edition

China: Please contact the customer service, Science Press, 16 Donghuangchenggen North Street, Beijing 100717, China. Tel: +86-10-64017032; E-mail: journal@mail.sciencep.com, or the local post office throughout China (domestic postcode: 2-580).

Outside China: Please order the journal from the Elsevier Customer Service Department at the Regional Sales Office nearest you.

Submission declaration

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis), that it is not under consideration for publication elsewhere. The submission should be approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out. If the manuscript accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder.

Submission declaration

Submission of the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis), that it is not under consideration for publication elsewhere. The publication should be approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out. If the manuscript accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder.

Editorial

Authors should submit manuscript online at <http://www.jesc.ac.cn>. In case of queries, please contact editorial office, Tel: +86-10-62920553, E-mail: jesc@263.net, jesc@rcees.ac.cn. Instruction to authors is available at <http://www.jesc.ac.cn>.

Journal of Environmental Sciences (Established in 1989)

Vol. 26 No. 12 2014

Supervised by	Chinese Academy of Sciences	Published by	Science Press, Beijing, China
Sponsored by	Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences		Elsevier Limited, The Netherlands
Edited by	Editorial Office of Journal of Environmental Sciences P. O. Box 2871, Beijing 100085, China Tel: 86-10-62920553; http://www.jesc.ac.cn E-mail: jesc@263.net , jesc@rcees.ac.cn	Distributed by	
		Domestic	Science Press, 16 Donghuangchenggen North Street, Beijing 100717, China Local Post Offices through China
		Foreign	Elsevier Limited http://www.elsevier.com/locate/jes
Editor-in-chief	Hongxiao Tang	Printed by	Beijing Beilin Printing House, 100083, China
CN 11-2629/X	Domestic postcode: 2-580	Domestic price per issue	RMB ¥ 110.00

ISSN 1001-0742



9 771001 074147